

CNS oxygen toxicity.

N. BITTERMAN

Technion, Israel Institute of Technology, Haifa 32000, Israel

INTRODUCTION

This short manuscript will briefly present our current knowledge about CNS oxygen toxicity: the clinical manifestations, descriptions and incidence of symptoms, and the time – duration relationship that defines the safety limits and risk factors leading to enhanced toxicity to hyperbaric oxygen. A very general outline will present suggested mechanisms underlying hyperoxia-induced seizures and strategies for protection against the seizures. Unfortunately, not much has been gained in our knowledge and understanding of CNS oxygen toxicity in recent years. Some concluding remarks will be given, as will some suggestions toward further research for increasing our knowledge about mechanisms, and better approaches for protection against the toxic effects of breathing oxygen at high partial pressures.

DESCRIPTION

There is no better description of CNS oxygen toxicity than the personal experience of Christian J. Lambertsen. “The convulsion is usually preceded but not always by the occurrence of localized muscular twitching, especially about the eyes, mouth and forehead. Small muscles of the hands may be involved and incoordination of diaphragm activity in respiration may occur. Eventually an abrupt spread of excitation occurs and the rigid "tonic" phase of the convulsions begins. Vigorous clonic contractions of the muscle groups of head and neck, trunk and limbs then occur becoming progressively less violent over 1 minute” (1). These convulsions are classified as generalized, tonic-clonic (grand mal) seizures (1). A typical electroencephalogram obtained during hyperbaric oxygen (HBO₂) seizures consists of spike and wave discharges (Figure 1) recorded in both hemispheres. They may start simultaneously or at random times in the cortical and subcortical areas (2, 3).

The hyperoxia-induced discharges are believed to be reversible, causing no residual neurological damage, and disappear upon reduction of the inspired oxygen partial pressure (1).




Fig. 1. Typical EEG from a rat exposed to 0.5 MPa oxygen. A) Air at atmospheric pressure. B) Appearance of typical spike and wave electrical discharges at 0.5 MPa oxygen. C) On return to atmospheric pressure. Calibration: 1 sec/50 V.

Hyperoxia-induced seizures are accepted as generalized, although several studies point toward local variations in

subcortical brain areas. This might contradict the general nature of the seizure's development and indicate a more localized primary focus for the initiation of the epileptic activity. Several studies demonstrate variations in cerebral blood flow in different brain areas on exposure to HBO₂ (4, 5, 6); others present regional changes in amino acids and ammonia levels (7), variation in lipid peroxide distribution (8), and distribution of antioxidant enzymes (9).

Early changes in cortical electrical activity have been described on exposure to HBO₂ several minutes prior to the full development of the electrical discharges (10, 11). Unfortunately, almost thirty years later despite extensive research, we have no real-time correlate of pre-seizure EEG activity that could serve for the prediction of CNS oxygen toxicity (12, 13, 14).

Symptoms

Few large-scale human studies describing the symptoms of CNS oxygen toxicity are available in the literature (15, 16, 17, 18, 19, 20). The list includes a group of minor symptoms, such as nausea, dizziness, abnormal sensations, headache, disorientation, lightheadedness and apprehension, which are difficult to define. More unambiguous symptoms appearing on exposure to HBO₂ are blurred vision, tunnel vision, and tinnitus. Signs include respiratory disturbances, eye twitching, twitching of lips, mouth, and forehead, and convulsions. There is no consistency in the appearance of symptoms and signs prior to the development of the seizures. Despite our prolonged experience with experimental and accidental CNS oxygen toxicity, there is no consensus about the frequency and incidence of the different symptoms.

Table 1 compares symptoms from about 550 human exposures to hyperbaric oxygen, compiled from those performed by Donald in 1947 (16), Leitch in 1984 (17), and three series of experiments conducted in Panama City by Butler and Thalmann (18, 19, 20). The symptoms of CNS oxygen toxicity are presented in the declining order of their incidences of appearance. As can be seen from the table, there are inconsistencies in the frequency of symptoms' appearance in the different HBO₂ series. This symptom variability reflects the importance of environmental and personal factors on the development of CNS oxygen toxicity.

Dosing

The most famous relationship between inspired oxygen partial pressures and the time for the appearance of symptoms of pulmonary and CNS oxygen toxicity (latent period) is the classic work of Lambertsen and his colleagues at the University of Pennsylvania (Figure 2) with exposures to HBO₂ in a dry hyperbaric chamber (21).

TABLE 1

Donald	Leitch	Butler and Thalmann (NEDU)		
1947	1984	1984	1986	1986
Convulsion	Convulsion	Light – Headedness	Nausea	Muscle twitching
Twitching of lips	Unconsciousness	Convulsion	Muscle twitching	Dizziness
Vertigo	Cyanosis	Tinnitus	Dizziness	Blurred vision
Nausea	Limb shaking	Apprehension	Tinnitus	Dysphoria
Respiratory Disturbances	Dizziness	Dysphoria	Dysphoria	Convulsion
Twitching of parts other than lips	Strenuous breathing	Blurred vision	Confusion	Aphasiaia
Sensations of abnormality	Auditory aura	Tunnel vision	Convulsion	Dyspnea
Visual Disturbances	Breathing disturbance	Disorientation	Decreased auditory Acuity	Paresthesias
Acoustic Hallucinations	Nausea	Lethargy	Aphasia	Nausea
Paraesthesiae	Dissociation	Dysphasia	Tingling	Lightheadedness
	Apnoea	Aphasia	Numbness	Air hunger
	Loud cry/groan	Eye twitching	Choking sensation	Tinnitus
	Malaise	Nystagmus	Amnesia	Confusion
	Headache/pulsation	Incoordination	Muscular rigidity	Muscular rigidity
	Apprehension		Lightheadedness	Irritability
	Amnesia		Poor concentration	Hypoacusis
	Facial twitch		Visual disturbances	Hyperacusis
	Lip tremor		Decreased mental alertness	Poor concetration
	Disorientation		Increased respiratory rate	Tunnel vision
N=388	N=35	N=28	N=33	N=59

Table 1. Symptoms of CNS oxygen toxicity presented in declining order of incidence. Data from Donald, 1947 (16), Leitch, 1984 (17), & Butler and Thalmann (18,19, 20).

Fig. 2. Pulmonary and neurological oxygen tolerance curves for continuous exposures of normal men to hyperbaric oxygen. Adapted from Lambertsen CJ, Clark JM, Gelfand R, et al, 1987 (21).

Figure 3 displays the depth- time limits for wet (immersed) oxygen diving (19), setting the lower limit for oxygen diving at about 15 msw (50 fsw) for a period of 10 minutes and allowing safe prolonged oxygen diving for about 4 hours at a fairly shallow depth of around 7 meters (20 fsw) (threshold depth). Various environmental and personal factors may modify sensitivity to CNS oxygen toxicity, thus shortening the duration of the latent period and lowering the threshold pressure for the development of seizures.






Fig. 3. Time–depth limits for oxygen diving. Data from Butler & Thalmann, 1986 (19).

Risk Factors

The exposure to hyperbaric oxygen in a wet environment increases sensitivity to CNS oxygen toxicity compared to exposure in a dry hyperbaric chamber (1, 16). Elevated concentrations of carbon dioxide (1, 14, 16, 22-25) and physical activity (exercise) dramatically decrease the duration of latent period for hyperoxia-induced seizures (1, 16, 23). The latent period for the appearance of electrical discharges in the EEG is significantly shorter in darkness than in light (26), suggesting the importance of visual input in the modulation of sensitivity to CNS oxygen toxicity. The risk for CNS oxygen toxicity is not determined solely by the partial pressure of the inspired oxygen, and even relatively low partial pressures of inert gases may contribute to hyperbaric-induced seizures (27). The increased sensitivity caused by inert gases could be explained by the involvement of free radical production (28). Circadian rhythm (29), various drugs, age (30), sex (31), interspecies differences and individual day to day variability (16) may contribute to the sensitivity to CNS oxygen toxicity (14). Figure 4 presents data from a unique study by Donald in 1947, in which he exposed a single diver to the same profile of HBO₂ for twenty times within a three month period, until the appearance of neurological symptoms of oxygen toxicity (16). As can be seen, there are large day to day variations in time duration of symptoms, suggesting that there is no fixed, personal, predetermined threshold of tolerance to oxygen toxicity. This finding supports that using the oxygen test as a screening tool for oxygen diver selection is ineffective.



In spite of the known day to day individual variability in tolerance to hyperoxic seizures, and the evidence concerning modulation of environmental and personal factors on the sensitivity to high oxygen pressures, it is surprising that we are still using tables based on statistical methods and predictive models. A more optimal approach would be personal monitoring of oxygen toxicity using a real-time predictive system for detecting the first signs of toxicity.

Fig. 4. Day to day variations in time duration until onset of CNS oxygen toxicity symptoms of a single diver. From Donald, 1947 (16).

Mechanisms

Although hyperoxic-induced seizures have been well described, the effects of HBO₂ on neural elements remain poorly understood. Studies of vertebrate nerves demonstrated decreased excitability and blockage of impulse conduction (32, 33), while increased axonal excitability and an increase in the membrane time constant were demonstrated in the isolated nervous system of the cockroach (34). The synaptic mechanisms seem to play an important role in the development of the HBO₂-induced seizures. Among the main electrophysiological findings is an increase in spontaneous synaptic transmitter release (35), a reduction in the inhibitory transmission and enhancement of evoked excitability activity at the neuromuscular junction of the lobster (36).

HBO₂ is known to affect most of the neurotransmitters and the neurotransmitter enzymes, such as GABA, acetylcholine, glutamate, dopamine, ammonia, norepinephrine, and aspartate (37-44). It also affects neuromodulators, such as nitric oxide (NO) (45). Membrane-bound active transport systems are also impaired on exposure to hyperbaric oxygen (14, 38, 39, 46, 47), with implications on neural activity. These studies were performed in different nerve preparations and synapse models. Some used direct physiological techniques, while others used indirect biochemical methods. They indicate that HBO₂ affects almost all neural elements. However, the primary neural target responsible for development of hyperoxic seizures is still unknown.

It is well accepted that reactive oxygen species (ROS) are increased on exposure to HBO₂, and overwhelming the body's normal antioxidant defenses may mediate the hyperoxic insult (48). An increase in free radical generation in the brain precedes HBO₂-induced convulsions, as demonstrated in brain extracts (49). H₂O₂ is elevated in various brain areas on exposure to HBO₂ (9, 50, 51), and there is a rise in free radical levels in the blood of humans exposed to HBO₂ (52). Nevertheless, a recent study done with awake animals using a microdialysis probe failed to detect an increase in hydroxyl radicals prior to, or during, HBO₂-induced convulsions (53). Currently, the available experimental data does not allow for us to decide whether ROS are the primary cause for hyperoxic seizures activity, or alternatively, that ROS are acting indirectly on neural elements and via second messengers, such as small inorganic molecules or proteins, to elicit the epileptic activity. A study using blood-brain barrier integrity as an index for injury in chronically EEG implanted rats revealed that CNS oxygen toxicity, at its

early stages (onset of electrical discharges), is not associated with the altered permeability of the cerebral microvessels (54).

Protection

Intermittent exposure to hyperbaric oxygen breathing with air breaks at the same pressure is a technical approach for increasing the total time of exposure to hyperoxia (14). This procedure is routinely used in the clinical setting. Various pharmacological strategies have been tested for postponing hyperoxic-induced seizures:

1. Cerebral vascular modulation. Exposure to HBO₂ results in cerebral vasoconstriction, leading to a decrease in CBF (4, 5, 1), which is later followed by cerebral vasodilatation. It has been suggested that the breakpoint in cerebral vasoconstriction is correlative with the development of hyperoxia-induced seizures (4). Therefore, any pharmacological agent that induces cerebral vasoconstriction may have the potential to protect, or at least postpone, the development of the seizures. Caffeine, a well known cerebral vasoconstrictor postpones hyperoxic seizures in a dose-related manner (55). Two nitric oxide synthase inhibitors, L-NAME and 7-nitroindazole (7-NI), significantly prolong the latent period before onset of seizures on exposure to hyperbaric oxygen or to a hypercapnic-hyperoxic stress (56, 57). This supports the involvement of the L-arginine-NO pathway in the pathophysiology of hyperoxia-induced seizures.
2. Neural activity modulation. On the basis of the clinical manifestations of hyperoxic seizures, several anti-epileptic drugs have been studied. A significant prolongation of the latent period before seizures was demonstrated using carbamazepine (Tegretol) (58). Vigabatrin (Gama vinyl GABA), an irreversible inhibitor of GABA transaminase, successfully protected against hyperoxic seizures in a dose-related manner for prolonged periods of 24 hours or more (59).
3. Enhancement of the antioxidant state. Extensive research had been directed toward defining agents that will protect against oxidative stress by increasing the potential for neutralizing or scavenging the ROS. The results up until now are not very promising. It is possible to find contradictory reports on various antioxidants' abilities to protect against CNS oxygen toxicity. For example, Puglia and Loeb (60) found that two of the most studied antioxidant enzymes, superoxide dismutase (SOD) and catalase (CAT), were protective, but both failed to exhibit protective effects in studies by Block, et al, (61). Others have shown that when entrapped in liposomes, SOD and CAT inhibit hyperoxia-induced seizures (62). No protection was found in transgenic animals over expressing SOD (63), and no effect was shown with knockout animals regarding pulmonary oxygen toxicity (64). Harabin, et al (65) have shown no correlation between the development of oxygen toxicity with intermittent exposures to hyperbaric oxygen and the levels of antioxidant enzymes.

Metal ion chelators, such as Deferoxamine, were not found to be protective against hyperoxic seizure (66). A variety of ROS scavengers and natural and synthetic antioxidants have been tested, showing different levels of protection in animal models (66-71). The Nitroxide spin traps Tempo and Tempol were found to be effective in preventing CNS oxygen toxicity and efficacy, correlated with their lipophilic properties (69, 70). Beta-carotene from natural origin (*Dunaliella bardawil*) was demonstrated to be among the most effective antioxidants after a one-week diet (71). The lack of dramatic protection against CNS oxygen toxicity by antioxidants is consistent with an assertion by two ROS pioneers in a recent review article. Gutteridge and

Halliwell write, “It was soon clear to many researchers that free radicals did not cause a plethora of diseases, neither were ‘spoonfuls’ of SOD or vitamins going to modify them, let alone cure them” (72). Deprivation of food or water prior to exposure to HBO₂ significantly prolonged the latent period to the onset of hyperoxia-induced seizures in a dose-related manner (73).

Further Research Areas

While trying to evaluate the current understanding of CNS oxygen toxicity, it is surprising that so much is unknown, or rather, that so little is known. Possible explanations for our limited knowledge in this field could be that HBO₂ is not a physiological signal, CNS oxygen toxicity is not a common disease, and for many years, HBO₂ diving was limited to combat military divers. Under such circumstances, not many research institutes or pharmaceutical companies undertake research or provide financial support for basic CNS oxygen toxicity studies. Many technical difficulties are imposed on experiments in hyperbaric conditions (74), whether simulated chamber exposures or deep-water diving (74). The candidate ROS that are suggested to mediate toxic effects have short half lives and are known to be site specific. This combination makes their real-time measurement a difficult task, especially in hyperbaric chambers. Oxygen toxicity is a multi-organ, multi-system toxicity. Therefore, it is difficult to separate the primary symptom from non-specific after effects. An additional reason for limited research may be the nature of the technological solutions available for avoiding CNS oxygen toxicity. They include using various mixtures with the precise control of oxygen partial pressure for diving and hyperbaric clinical use.

CONCLUDING REMARKS

1. An effort should be directed at establishing the importance and utility of hyperoxia-induced seizures as an experimental model of generalized (grand mal) epilepsy (75). Hyperoxia-induced seizures can serve as an excellent epileptic model since the epileptic agent disappears immediately on reduction of the oxygen pressure, and its toxic manifestations are believed to be reversible. Collaborations among scientists from the basic neurological research (especially the mechanisms of epilepsy) will increase our knowledge and improve our ability to use innovative strategies for the prevention of CNS oxygen toxicity.
2. We must continue to expand research efforts on the basic mechanism of oxidative stress and the role of second messengers in the pathophysiology of CNS oxygen toxicity. This approach will increase the need for more sophisticated methods of multimodal monitoring of physiological and biochemical parameters, such as mentioned by Rogatsky, Shifrin, and Mayevsky (76). A future protection strategy should consider the need for a continuous inflow of antioxidants or scavengers to neutralize the ROS or avoid its development in real time, during the entire period of exposure to HBO₂. That is, a patch that releases a continuous flow or an implanted biological generator of ROS scavengers.
3. An extensive effort must be given to having a personalized, real-time profile for any HBO₂ diver or patient, based on his personal, daily-specific sensitivity to HBO₂, instead of using tables of the time–duration relationship. In our technological era, this can be done with miniature sensors and wireless technology, which will follow the physiological parameters and pre-seizure modifications. The future vision for oxygen diving will be in a tailored CNS oxygen toxicity monitor, which will assist in allowing each person to have an optimal level of hyperbaric oxygen without being exposed to its toxicity.

REFERENCES

1. Lambertsen CJ. Effects of oxygen at high partial pressure. In: Fenn WO, Rahn H., eds. Handbook of Physiology. Respiration. Bethesda, MD: *Am. Physiol. Soc.*, Sec 3, Vol 2, 1965: 1027-1046.
2. Harel D, Kerem D, and Lavy S. The influence of high oxygen pressure on the electrical activity of the brain. *Electroenceph Clin Neurophysiol* 1969; 26:310-317.
3. Rucci FS, Givetti HL, and La Rocca H. Changes in electrical activity of the cerebral cortex and of some subcortical centers in hyperbaric oxygen. *Electroenceph Clin Neurophysiol* 1967; 22:231-238.
4. Torbati D, Parolla D, and Lavy S. Blood flow in rat brain during exposure to high oxygen pressure. *Aviat Space Environ Med* 1978; 49:963-967.
5. Bergo GW and Tyssebotn I. Cerebral blood flow distribution during exposure to 5 bar oxygen in awake rats. *Undersea Biomed Res* 1992; 19:339-354.
6. Demchenko IT, Boso AE, Oneill TJ, Bennett PB, and Piantadosi CA. Nitric oxide and cerebral blood flow responses in hyperbaric oxygen *J Appl Physiol* 2000; 88:381-1389.
7. Mialon P, Gibey R, Bigot JC, and Barthelemy L. Changes in striatal and cortical amino acid and ammonia levels of rat brain after one hyperbaric oxygen-induced seizure. *Aviat Space Environ Med* 1992; 63:287-291.
8. Noda Y, McGreer PL, and McGreer EG. Lipid peroxide distribution in brain and the effect of hyperbaric oxygen. *J Neurochem* 1983; 40:1329-1332.
9. Piantadosi CA and Tatro LG. Regional H₂O₂ concentration in rat brain after hyperoxic convulsions. *J Appl Physiol* 1990; 69:1761-1766.
10. Rada-Bitterman N, Conforti N, Harel D, and Lavy S. Analysis of pre-seizure EEG changes in rats during hyperbaric oxygenation. *Exp Neurol* 1975; 46:9-19.
11. Torbati D, Simon AJ, and Ranade A. Frequency analysis of EEG in rats during the preconvulsive period of O₂ poisoning. *Aviat Space Environ Med* 1981; 52:598-603.
12. Visser GH, Van Hulst RA, Wieneke GH, and Van Huffelen AC. The contribution of conventional and quantitative electroencephalography during monitoring of exposure to hyperbaric oxygen. *Undersea Hyperb Med* 1996; 23:91-98.
13. Geva AB and Kerem DH. Forecasting generalized epileptic seizure from the EEG signal by wavelet analysis and dynamic unsupervised fuzzy clustering. *IEEE Trans Biomed Eng* 1998; 45:1205-1216.
14. Clark JM. Oxygen Toxicity. In: Bennett PB, Elliott DH, eds. The physiology and medicine of diving. London, Saunders, 1993: 121-169.
15. Donald KW. Oxygen poisoning in man. *Brit Med J* 1947; 1:667-712, 712-717.
16. Donald KW. Oxygen and the diver. Harley Swan, UK. The SPA Ltd., 1992.
17. Leitch DR. Cerebral oxygen toxicity during immersion and surface supported diving. INM TECH Memo 12/84. 1984.
18. Butler FK and Thalmann ED. CNS oxygen toxicity in closed-circuit scuba diving. In: Bachrach AJ, Matzen MM, eds. Proc Eighth Symp Underwater Physiol. Bethesda, Md: Undersea Medical Society.1984: 15-30.
19. Butler FK and Thalmann ED. Central nervous system oxygen toxicity in closed circuit scuba divers II. *Undersea Biomed Res* 1986; 13:193-223.
20. Butler FK. Central nervous system oxygen toxicity in closed circuit scuba divers III. US Navy Experimental Diving Unit Report No.5-86 .1986.
21. Lambertsen CJ, Clark JM, Gelfand R, et al. Definition of tolerance to continuous hyperoxia in man: an abstract report of predictive studies V. In: Bove AA, Bachrach AJ, Greenbaum LJ, eds. Proc 9th International Symposium on Underwater and Hyperbaric Physiology. Undersea and Hyperbaric Society, Bethesda, MD. 1987: 717-735.
22. Clark JM. Effects of acute and chronic hypercapnia on oxygen tolerance in rats. *J Appl Physiol: Resp Environ Exercise Physiol* 1981; 50:1036-1044.
23. Kaffka M, Bitterman N, and Kerem D. The enhancement of oxygen toxicity by exercise & CO₂: An animal model. In: Abstracts of the XII Meeting of the European Undersea Biomedical Society, Rotterdam, Holland, August 27-30, 1986.
24. Arieli R and Ertracht O. Latency to CNS oxygen toxicity in rats as a function of PCO₂ and PO₂. *Eur J Appl Physiol Occup Physiol* 1999; 80:598-603.
25. Bitterman N and Bitterman H. CNS oxygen toxicity: Diverse Protection of antiepileptic and vasoactive agents during exposure to HBO₂ or HBO₂+CO₂. *Undersea Hyperb Med* 1999; 26 (Suppl): 39.

26. Bitterman N, Melamed Y, and Perlman I. CNS Oxygen toxicity in the rat: The role of ambient illumination. *Undersea Biomed Res* 1986; 13:19-25.
27. Bitterman N, Laor A, and Melamed Y. CNS oxygen toxicity in gas mixture breathing. *Undersea Biomed Res* 1987; 14:477-483.
28. Thom SR. Inert gas enhancement of superoxide radical production. *Arch Biochem Biophys* 1992; 295:391-396.
29. Hof DG, Dexter JD, and Mengel CE. Effect of circadian rhythm on CNS oxygen toxicity. *Aerospace Med* 1971; 42:1293-1296.
30. Torley LW and Weiss HS. Effect of age and magnesium ions on oxygen toxicity in the neonate chicken. *Undersea Biomed Res* 1975; 2:223-227.
31. Troy SS and Ford DH. Hormonal protection of rats breathing oxygen at high pressure. *Acta Neurol Scand* 1972; 48:231-242.
32. Cymerman A and Gottlieb SF. Effects of increased oxygen tensions on bioelectric properties of frog sciatic nerve. *Aerosp Med* 1970; 41:36-39.
33. Perot PL and Stein SN. Conduction block in mammalian nerve produced by oxygen at high pressure. *Am J Physiol* 1959; 197:1243-1246.
34. Bitterman N and Grossman Y. Increased axonal excitability during exposure to hyperbaric oxygen. *Can J Physiol Pharmacol* 1987; 65:2044-2047.
35. Colton JS and Colton CA. Effect of oxygen at high pressure on spontaneous transmitter release. *Am J Physiol* 1978; 238:C233-C237.
36. Colton JS and Colton CA. The action of oxygen and oxygen at high pressures on inhibitory transmission. *Brain Res* 1986; 364:151-158.
37. Tunnicliff G, Urton M, and Wood JD. Susceptibility of chick brain L-glutamic acid decarboxylase and other neurotransmitter enzymes to hyperbaric oxygen in vitro. *Biochem Pharmacol* 1973; 22:501-505.
38. Wood JD, Watson WJ and Murray GW. Correlation between decrease in brain γ -aminobutyric acid levels and susceptibility to convulsions induced by hyperbaric oxygen. *J Neurochem* 1969; 16:281-287.
39. Zirkle LG, Mengel CE, Horton BD, and Duffy EJ. Studies of oxygen toxicity in the central nervous system. *Aerosp Med* 1965; 36:1027-1032.
40. Davis HC and Davis RE. Biochemical aspects of oxygen poisoning. In: Fenn WO, Rahn H., eds. Handbook of Physiology. Respiration. Bethesda, MD: *Am. Physiol. Soc.*, Sec 3, Vol 2, 1965: 1047-1058.
41. Haugaard N. Cellular mechanisms of oxygen toxicity. *Physiol Rev* 1968; 48:11-373.
42. Fisher DB and Kaufman S. The inhibition of phenylalanine and tyrosine hydroxylases by high oxygen levels. *J Neurochemistry* 1972; 19:1359-1365.
43. Mialon P, et al. Amino acids and ammonia in the cerebral cortex, the corpus striatum and the brain stem of the mouse prior to the onset and after a seizure induced by hyperbaric oxygen. *Brain Res* 1995; 676:352-357.
44. Huggins AK and Nelson DR. The effects of hyperbaric oxygenation on the levels of 5-hydroxytryptamine, noradrenaline, dopamine and free amino acids in whole mouse brain. *J Neurochem* 1975; 25:117-121.
45. Demchenko IT, Boso AE, Whorton AR, and Piantadosi CA. Nitric oxide production is enhanced in rat brain before oxygen-induced convulsions. *Brain Res* 2001; 917:253-261.
46. Gottlieb SF and Cymerman A. Effects of increased oxygen tension on sodium active transport through frog skin. *Aerosp Med* 1970; 41:661-665.
47. Kovachich GB, Mishra OP, and Clark JM. Depression of cortical Na, K-ATPase activity on rats exposed to hyperbaric oxygen. *Brain Res* 1981; 206:229-232.
48. Gershman R, Gilbert DL, Nye SW, Dwyer P, and Fenn WO. Oxygen poisoning and X-irradiation: A mechanism in common. *Science* 1954; 119:623-626.
49. Torbati D, Church DF, Keller JM, and Pryor WA. Free radical generation in the brain precedes hyperbaric oxygen induced convulsions. *Free Radical Biol Med* 1992; 13:101-106.
50. Jerrett SA, Jefferson D, and Mengel CE. Seizures, H₂O₂ formation and lipid peroxides in brain during exposures to oxygen under high pressure. *Aerosp Med* 1973; 44:40-44.
51. Yusa T, Beckman JS, Crapo JD, and Freeman BA. Hyperoxia increase H₂O₂ production by brain in vivo. *J Appl Physiol* 1987; 63:353-358.
52. Narkowicz CK, Vial JH, and McCartney PW. Hyperbaric oxygen therapy increases free radical levels in the blood of human. *Free Radic Res Commun* 1993; 19:71-80.
53. Amiridze N, Dang Y, and Brown OR. Hydroxyl radicals detected via brain microdialysis in rats breathing air and during hyperbaric oxygen convulsions. *Redox Rep* 1999; 4:165-170.
54. Gross B, Bitterman N, Levanon D, Nir I, and Harel D. Horseradish peroxidase as a marker of blood-brain barrier integrity in oxygen toxicity in the central nervous system. *Exptl Neurol* 1986; 93:471-480.

55. Bitterman N and Schaal S. Caffeine attenuates CNS oxygen toxicity in rats. *Brain Res* 1995; 696:250-253.
56. Bitterman N, Bitterman H. L-arginine-NO pathway and CNS oxygen toxicity. *J Appl Physiol* 1997; 84: 1633-1638.
57. Zhang J, Su Y, Oury TD, and Piantadosi A. Cerebral amino acid, norepinephrine, and nitric oxide metabolism in CNS oxygen toxicity. *Brain Res* 1993; 606:56-62.
58. Reshef A, Bitterman N, and Kerem D. The effect of Carbamazepine and Ethosuximide on hyperoxic seizures. *Epilepsy Res.* 1991; 8:117-121.
59. Tzuk-Shina T, Bitterman N, and Harel D. The effect of Vigabatrin on central nervous system oxygen toxicity in rats. *Euro J Pharmacol.* 1991; 202:171-175.
60. Puglia CD and Loeb GA. Influence of rat brain superoxide dismutase inhibition by diethyldithiocarbamate upon the rate of development of central nervous system oxygen toxicity. *Toxicol Appl Pharmacol* 1984; 75:258-264.
61. Block ER. Effect of superoxide dismutase and succinate on the development of hyperbaric oxygen toxicity. *Aviat Space Environ Med* 1977; 48:645-648.
62. Yusa T, Crapo JD, and Freeman BA. Liposome-mediated augmentation of brain SOD and catalase inhibits CNS oxygen toxicity. *J Appl Physiol* 1984; 57:1674-1681.
63. Oury TD, Ho YS, Piantadosi CA, and Crapo JD. Extracellular superoxide dismutase, nitric oxide, and central nervous system in oxygen toxicity. *Proc Natl Acad Sci* 1992; 89:9715-9719.
64. Tsan MF, White JE, Caska B, Epstein CJ, and Lee CY. Susceptibility of heterozygous MnSOD gene-knockout mice to oxygen toxicity. *Am J Respir Cell Mol Biol* 1998; 19:114- 120.
65. Harabin AL, Survanshi SS, Weathersby PK, Hays JR, and Homer LD. The modulation of oxygen toxicity by intermittent exposure. *Toxicol Appl Pharmacol* 1988; 93:298-311.
66. Lin Y and Jamieson D. Effects of antioxidants on oxygen in vivo and lipid peroxidation in vitro. *Pharmacol Toxicol* 1992; 70:271-277.
67. Boadi WY, Thaire L, Kerem D, and Yannai S. Effects of dietary supplementation with vitamin E, riboflavin and selenium on central nervous system oxygen toxicity. *Pharmacol Toxicol* 1991; 68:77-82.
68. Jamieson D and Van den Brenk HAS. The effect of antioxidants on high pressure oxygen toxicity. *Biochem Pharmacol* 1964; 13:159-164.
69. Bitterman N and Samuni A. Nitroxide stable radicals protect against hyperoxic induced seizures in rats. *Undersea Hyperbaric Med* 1995;22 (Suppl): 47-
70. Bitterman N and Samuni A. Albumin conjugated nitroxide protects against CNS oxygen toxicity in rats. In: Proceedings of the Annual Meeting of the E.U.B.S. on Diving and Hyperbaric Medicine, Stockholm, Sweden. August 12-15, 1998.
71. Bitterman N, Melamed Y, and Ben-Amotz A. Beta-carotene and CNS oxygen toxicity. *J Appl Physiol* 1994; 76:1073-1076.
72. Gutteridge JM and Halliwell B. Free radicals and antioxidants in the year 2000. A historical look to the future. *Ann N Y Acad Sci* 2000; 899:136-147.
73. Bitterman N, Skapa E, and Gutterman A. Starvation and dehydration attenuate CNS oxygen toxicity in rats. *Brain Res* 1997; 761:146-150.
74. Dean JB and Mulky DK. Continuous intracellular recording from mammalian neurons exposed to hyperbaric helium, oxygen, or air. *J Appl Physiol* 2000; 89:807-822.
75. Wood JD. Systemic oxygen derangements. In: Purpura DP, Penry JK, Tower D, Woodbury DM, Walter R. eds. Experimental models of epilepsy. Raven Press, 1972: 459-477.
76. Rogatsky GG, Shifrin EG, and Mayevsky A. Physiological and biochemical monitoring during hyperbaric oxygenation. A review. *Undersea Hyperb Med* 1999; 26:111-122.